


ADA 035572

ORD. 12640. 16 Dr
✓ 

RESEARCH REPORT

DDC
RECEIVED
FEB 14 1971
REGISTRY
D

NOT AVAILABLE TO DDC DOES NOT
WARRANT FULLY LEGIBLE PRODUCTION

Industrial & Systems
Engineering Department
University of Florida
Gainesville, FL. 32611

DISTRIBUTION STATEMENT A
Approved for public release
Distribution Unlimited

CLASSIFIED BY	
DTIC	Uncl. Status <input checked="" type="checkbox"/>
DDI	Uncl. Status <input type="checkbox"/>
UNCLASSIFIED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODE	
Dist.	AVAIL. and/or SPECIAL
A	


The Incomplete Dirichlet's Multiple Integral

Research Report No. 76-24

by

B. D. Sivazlian

November, 1976



Department of Industrial and Systems Engineering
University of Florida
Gainesville, Florida 32611

APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED

This research was supported in part by the Army Research Office,
Triangle Park, NC, under contract number DAHCO4-75-0150.

THE FINDINGS OF THIS REPORT ARE NOT TO BE CONSTRUED AS AN OFFICIAL
DEPARTMENT OF THE ARMY POSITION, UNLESS SO DESIGNATED BY OTHER
AUTHORIZED DOCUMENTS.

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 76-24	2. SOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER (9) Research rept.
4. TITLE (and Subtitle) (6) The Incomplete Dirichlet's Multiple Integral		5. TYPE OF REPORT & PERIOD COVERED Technical
7. AUTHOR(s) (10) B. D. Sivazlian	(14) RA-76-24	6. PERFORMING ORG. REPORT NUMBER 76-24
	(15)	8. CONTRACT OR GRANT NUMBER(s) DAHC04-75-0150 new
9. PERFORMING ORGANIZATION NAME AND ADDRESS Industrial and Systems Engineering University of Florida Gainesville, FL 32611		10. PROGRAM ELEMENT, PROJECT, TASK AREA & REPORT NUMBER (16) 20661102A14D Rsch in & Appl of Applied Math.
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Triangle Park, NC 27709		12. REPORT DATE (17) November, 1976
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 27 (13) 29 p.
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited		(18) ARO (19) 12640-16-M
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) N/A		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Multiple integrals Incomplete beta functions Multivariate statistical distributions Dirichlet's distribution Beta distribution		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) It is shown that this integral is reducible to a single integral involving products of incomplete beta functions as a multiplicative constant. The results are generalized and extended to other multiple integrals. Finally, as an application a class of multivariate type distribution defined by a functional form is partially investigated.		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

404399
698

Table of Contents

ABSTRACT	1
SECTIONS	
1. Introduction.	1
2. The Incomplete Dirichlet's Multiple Integral.	2
3. Some Generalizations.	6
4. Extensions.	8
5. A Still Further Extension	11
6. Applications to Statistics.	12
REFERENCES	22

Abstract

The incomplete Dirichlet's multiple integral is defined as

$$\int \int \cdots \int_R f(t_1 + t_2 + \cdots + t_n) t_1^{\alpha_1-1} t_2^{\alpha_2-1} \cdots t_n^{\alpha_n-1} dt_1 dt_2 \cdots dt_n$$

where $R = \{(t_1, t_2, \dots, t_n): t_i \geq 0, \sum_{i=1}^n t_i \leq 1, t_{i+1} \leq a_i \sum_{j=1}^i t_j; i = 1, 2, \dots, n-1\}$

It is shown that this integral is reducible to a single integral involving products of incomplete beta functions as a multiplicative constant. The results are generalized and extended to other multiple integrals. Finally, as an application a class of multivariate type distribution defined by a functional form is partially investigated.

1. Introduction

If $f(\cdot)$ is a continuous function of some real non-negative argument, and $\alpha_i > 0$, $i = 1, 2, \dots, n$, then the Dirichlet's multiple integral

$$I_n = \int \int \dots \int_{t_1+t_2+\dots+t_n \leq 1} f(t_1 + t_2 + \dots + t_n) t_1^{\alpha_1-1} t_2^{\alpha_2-1} \dots t_n^{\alpha_n-1} dt_1 dt_2 \dots dt_n \quad (1)$$

may be expressed as the single integral [5], [18]

$$I_n = \frac{\Gamma(\alpha_1)\Gamma(\alpha_2) \dots \Gamma(\alpha_n)}{\Gamma(\alpha_1 + \alpha_2 + \dots + \alpha_n)} \int_0^1 f(\tau) \tau^{(\alpha_1+\alpha_2+\dots+\alpha_n)-1} d\tau$$

Alternatively, we may express the multiplicative quotient involving gamma functions in terms of products of beta functions and write

$$I_n = B(\alpha_2, \alpha_1) B(\alpha_3, \alpha_2 + \alpha_1) \dots B(\alpha_n, \alpha_{n-1} + \alpha_{n-2} + \dots + \alpha_2 + \alpha_1) \int_0^1 f(\tau) \tau^{(\alpha_1+\alpha_2+\dots+\alpha_n)-1} d\tau \quad (2)$$

where in general

$$B(l, m) = \frac{\Gamma(l)\Gamma(m)}{\Gamma(l+m)}$$

An extension of (1) was proposed by Sivazlian [16] where in (1) the argument of the function $f(\cdot)$ was replaced by various partial sums of the variables t_1, t_2, \dots, t_n . Further extensions in the integrand form of (1) were proposed by Sivazlian [17] and Klamkin [13]. Klamkin also suggested an interesting extension of (1) in yet another direction, namely by changing the region of integration to $\{(t_1, t_2, \dots, t_n): t_i \geq 0, \sum_{i=1}^r t_i \leq t, \sum_{i=r+1}^n t_i \leq t'\}$, $t, t' > 0$. The multiple integral is then reducible to a double integral again involving quotients of gamma functions as its multiplicative constant.

In the present work, we propose still another region R defined for $0 < a_i \leq \infty$, $i = 1, 2, \dots, n$ as the simplex

$$R = \{(t_1, t_2, \dots, t_n): t_i \geq 0, \sum_{i=1}^n t_i \leq 1, t_{i+1} \leq a_i \sum_{j=1}^i t_j; \\ i = 1, 2, \dots, n-1\} \quad (3)$$

and define the incomplete Dirichlet's multiple integral as

$$I_n = \int \int \dots \int_R f(t_1 + t_2 + \dots + t_n) t_1^{\alpha_1-1} t_2^{\alpha_2-1} \dots t_n^{\alpha_n-1} dt_1 dt_2 \dots dt_n \quad (4)$$

This integral is not reducible to any of the previous forms investigated.

However, in the very special case when $a_i = \infty$, the Liouville's extension of Dirichlet's integral, that is form (1) is recovered. In general, we shall show that the proposed multiple integral (4) has the same format as (2) except that the multiplicative constants outside the single integral are products of incomplete beta functions. We shall show that $I_n = kI_n$, where k is the product of $(n-1)$ incomplete beta function ratio.

2. The Incomplete Dirichlet's Multiple Integral

To reduce (4), we first make the change in variable $t_i = x_i^2$, $i = 1, 2, \dots, n$. The Jacobian of the transformation is clearly $2^n x_1 x_2 \dots x_n$. The new region of integration \hat{R} is defined by the set of points

$$\hat{R} \equiv \{(x_1, x_2, \dots, x_n): x_i \geq 0, \sum_{i=1}^n x_i^2 \leq 1, x_{i+1} \leq \sqrt{a_i} \sqrt{x_1^2 + x_2^2 + \dots + x_i^2} \\ \text{for } i = 1, 2, \dots, n-1\}$$

Hence

$$I_n = 2^n \int \int \dots \int_{\hat{R}} f(x_1^2 + x_2^2 + \dots + x_n^2) x_1^{2\alpha_1-1} x_2^{2\alpha_2-1} \dots x_n^{2\alpha_n-1} dx_1 dx_2 \dots dx_n$$

We now utilize a second change in variable which is the elegant transformation into generalized n -dimensional spherical coordinates proposed by Schlafli in 1855 [4], also by Clare in 1881 [5], and used often in statistics (e.g. see [12]). Let then

$$\begin{aligned}
x_1 &= r \cos \theta_{n-1} \cos \theta_{n-2} \cdots \cos \theta_3 \cos \theta_2 \cos \theta_1 \\
x_2 &= r \cos \theta_{n-1} \cos \theta_{n-2} \cdots \cos \theta_3 \cos \theta_2 \sin \theta_1 \\
x_3 &= r \cos \theta_{n-1} \cos \theta_{n-2} \cdots \cos \theta_3 \sin \theta_2 \\
&\dots \\
x_{n-1} &= r \cos \theta_{n-1} \sin \theta_{n-2} \\
x_n &= r \sin \theta_{n-1}
\end{aligned}$$

The Jacobian of the transformation is

$$r^{n-1} (\cos \theta_{n-1})^{n-2} (\cos \theta_{n-2})^{n-3} \cdots (\cos \theta_2)$$

To determine the new region of integration \tilde{R} , we note that the ball

$0 \leq \sum_{i=1}^n x_i^2 \leq 1$ is mapped into the set of points $0 \leq r \leq 1$. Also, for $i = 1, 2, \dots, n-1$, the set of points defined by

$$0 \leq x_{i+1} \leq \sqrt{a_i} \sqrt{x_1^2 + x_2^2 + \cdots + x_i^2}$$

is mapped into the set of points

$$\begin{aligned}
0 &\leq r \cos \theta_{n-1} \cos \theta_{n-2} \cdots \cos \theta_{i+2} \cos \theta_{i+1} \sin \theta_i \\
&\leq \sqrt{a_i} r \cos \theta_{n-1} \cos \theta_{n-2} \cdots \cos \theta_{i+1} \cos \theta_i
\end{aligned}$$

$$\text{or } 0 \leq \tan \theta_i \leq \sqrt{a_i}$$

$$\text{or } 0 \leq \theta_i \leq \tan^{-1} \sqrt{a_i}$$

It thus follows that \hat{R} is mapped into \tilde{R} where

$$\begin{aligned}
\tilde{R} &= \{(r, \theta_1, \theta_2, \dots, \theta_{n-1}) : 0 \leq r \leq 1, 0 \leq \theta_i \leq \tan^{-1} \sqrt{a_i} \\
&\quad \text{for } i = 1, 2, \dots, n-1\}
\end{aligned}$$

The expression for \tilde{I} can then be written as

$$\begin{aligned}
\tilde{I} &= 2^n \int \int \cdots \int_{\tilde{R}} f(r^2) (r \cos \theta_{n-1} \cos \theta_{n-2} \cdots \cos \theta_2 \cos \theta_1)^{2\alpha_1-1} \\
&\quad (r \cos \theta_{n-1} \cos \theta_{n-2} \cdots \cos \theta_2 \sin \theta_1)^{2\alpha_2-1} \\
&\quad \cdots (r \cos \theta_{n-1} \sin \theta_{n-2})^{2\alpha_{n-1}-1} (r \sin \theta_{n-1})^{2\alpha_n-1} \\
&\quad [r^{n-1} (\cos \theta_{n-1})^{n-2} (\cos \theta_{n-2})^{n-3} \cdots (\cos \theta_2)] dr d\theta_1 d\theta_2 \cdots d\theta_n \\
&= 2^n \left[\int_0^1 r^{2(\alpha_1+\alpha_2+\cdots+\alpha_n)-1} f(r^2) dr \right]
\end{aligned}$$

$$\begin{aligned}
& \left[\int_0^{\tan^{-1}\sqrt{a_1}} (\sin \theta_1)^{2\alpha_2-1} (\cos \theta_1)^{2\alpha_1-1} d\theta_1 \right] \\
& \left[\int_0^{\tan^{-1}\sqrt{a_2}} (\sin \theta_2)^{2\alpha_3-1} (\cos \theta_2)^{2(\alpha_1+\alpha_2)-1} d\theta_2 \right] \\
& \dots \\
& \left[\int_0^{\tan^{-1}\sqrt{a_{n-2}}} (\sin \theta_{n-2})^{2\alpha_{n-1}-1} (\cos \theta_{n-2})^{2(\alpha_1+\alpha_2+\dots+\alpha_{n-2})-1} d\theta_{n-2} \right] \\
& \left[\int_0^{\tan^{-1}\sqrt{a_{n-1}}} (\sin \theta_{n-1})^{2\alpha_n-1} (\cos \theta_{n-1})^{2(\alpha_1+\alpha_2+\dots+\alpha_{n-1})-1} d\theta_{n-1} \right]
\end{aligned}$$

Making the change in variable $\tau = r^2$ and $u_i = \sin^2 \theta_i$, $i = 1, 2, \dots, n-1$, we obtain

$$\begin{aligned}
I_n &= \left[\int_0^{\frac{a_1}{1+a_1}} u_1^{\alpha_2-1} (1-u_1)^{\alpha_1-1} du_1 \right] \left[\int_0^{\frac{a_2}{1+a_2}} u_2^{\alpha_3-1} (1-u_2)^{\alpha_2+\alpha_1-1} du_2 \right] \\
&\dots \left[\int_0^{\frac{a_{n-1}}{1+a_{n-1}}} u_{n-1}^{\alpha_n-1} (1-u_{n-1})^{\alpha_{n-1}+\alpha_{n-2}+\dots+\alpha_2+\alpha_1-1} du_{n-1} \right] \\
&\quad \left[\int_0^1 f(\tau) \tau^{\alpha_1+\alpha_2+\dots+\alpha_{n-1}} d\tau \right]
\end{aligned}$$

Using the definition of the incomplete beta function

$$B_x(\alpha, \beta) = \int_0^x y^{\alpha-1} (1-y)^{\beta-1} dy$$

we obtain the following expression for I_n :

$$\begin{aligned}
I_n &= \left[B_{\frac{a_1}{1+a_1}}(\alpha_2, \alpha_1) \right] \left[B_{\frac{a_2}{1+a_2}}(\alpha_3, \alpha_2 + \alpha_1) \right] \dots \left[B_{\frac{a_{n-1}}{1+a_{n-1}}}(\alpha_n, \alpha_{n-1} + \right. \\
&\quad \left. \alpha_{n-2} + \dots + \alpha_1) \right] \\
&\quad \cdot \int_0^1 f(\tau) \tau^{\alpha_1+\alpha_2+\dots+\alpha_{n-1}} d\tau
\end{aligned} \tag{5}$$

Finally, if $I_x(a, b) = B_x(a, b)/B(a, b)$ is the incomplete beta function ratio, then clearly:

$$I_n/I_n = \prod_{i=1}^{n-1} I_{\frac{a_i}{1+a_i}}(\alpha_i, \alpha_{i-1} + \alpha_{i-2} + \dots + \alpha_1)$$

which is independent of the function $f(\cdot)$. In the special case when $a_i = \infty$ for all $i = 1, 2, \dots, n - 1$, the value of the previous expression equals 1.

3. Some Generalizations

We consider two generalizations of particular interest although other classes of multiple integrals may be treated in the same fashion. First consider the multiple integral

$$I_n(t) = \int \int \dots \int_S f(u_1 + u_2 + \dots + u_n) u_1^{\alpha_1-1} u_2^{\alpha_2-1} \dots u_n^{\alpha_n-1} du_1 du_2 \dots du_n \quad (6)$$

where S is the simplex defined by the set of points

$$S = \{(u_1, u_2, \dots, u_n) : u_i \geq 0, \sum_{i=1}^n u_i \leq t, u_{i+1} \leq a_i \sum_{j=1}^i u_j; \\ i = 1, 2, \dots, n-1\}$$

Making the change in variables $u_i = t t_i$, $i = 1, 2, \dots, n$, we obtain

$$I_n(t) = t^{\alpha_1 + \alpha_2 + \dots + \alpha_n} \int \int \dots \int_R f[t(t_1 + t_2 + \dots + t_n)] \\ t_1^{\alpha_1-1} t_2^{\alpha_2-1} \dots t_n^{\alpha_n-1} dt_1 dt_2 \dots dt_n$$

where R is the same simplex as (3). Using (5) and after making an obvious change in variable we obtain

$$I_n(t) = [B_{\frac{a_1}{1+a_1}}(\alpha_2, \alpha_1)] [B_{\frac{a_2}{1+a_2}}(\alpha_3, \alpha_2 + \alpha_1)] \dots [B_{\frac{a_{n-1}}{1+a_{n-1}}}(\alpha_n, \alpha_{n-1} + \alpha_{n-2} + \dots + \alpha_1)] \int_0^t f(\tau) \tau^{\alpha_1 + \alpha_2 + \dots + \alpha_n - 1} d\tau \quad (7)$$

(7) is of the same form as (5) except for the upper limit of the integral.

We next consider the integral

$$K_n(t) = \int \int \dots \int_T f\left[\left(\frac{v_1}{q_1}\right)^{\beta_1} + \left(\frac{v_2}{q_2}\right)^{\beta_2} + \dots + \left(\frac{v_n}{q_n}\right)^{\beta_n}\right] \cdot \\ v_1^{\alpha_1-1} v_2^{\alpha_2-1} \dots v_n^{\alpha_n-1} dv_1 dv_2 \dots dv_n \quad (8)$$

where T is the region of integration defined by the set of points

$$T = \{(v_1, v_2, \dots, v_n) : v_i \geq 0, \sum_{i=1}^n \left(\frac{v_i}{q_i}\right)^{\beta_i} \leq t, \left(\frac{v_{i+1}}{q_{i+1}}\right)^{\beta_{i+1}} \leq a_i \sum_{j=1}^i \left(\frac{v_j}{q_j}\right)^{\beta_j}; i = 1, 2, \dots, n\}$$

under the assumptions that $\beta_i > 0$, $\alpha_i > 0$, $q_i > 0$ for $i = 1, 2, \dots, n$. Here,

making the change in variable $\left(\frac{v_i}{q_i}\right)^{\beta_i} = u_i$, $i = 1, 2, \dots, n$, we obtain

$$K_n(t) = \frac{q_1^{\alpha_1} q_2^{\alpha_2} \dots q_n^{\alpha_n}}{\beta_1 \beta_2 \dots \beta_n} \int \int \dots \int_S f(u_1 + u_2 + \dots + u_n) u_1^{\frac{\alpha_1}{\beta_1} - 1} u_2^{\frac{\alpha_2}{\beta_2} - 1} \dots u_n^{\frac{\alpha_n}{\beta_n} - 1} du_1 du_2 \dots du_n$$

which is of the same form as (6). Hence, using (7) we obtain

$$K_n(t) = \frac{q_1^{\alpha_1} q_2^{\alpha_2} \dots q_n^{\alpha_n}}{\beta_1 \beta_2 \dots \beta_n} [B_{\frac{\alpha_1}{\beta_1}} \left(\frac{\alpha_2}{\beta_2}, \frac{\alpha_1}{\beta_1}\right) [B_{\frac{\alpha_2}{\beta_2}} \left(\frac{\alpha_3}{\beta_3}, \frac{\alpha_2}{\beta_2} + \frac{\alpha_1}{\beta_1}\right) \dots [B_{\frac{\alpha_{n-1}}{\beta_{n-1}}} \left(\frac{\alpha_n}{\beta_n}, \frac{\alpha_{n-1}}{\beta_{n-1}} + \frac{\alpha_{n-2}}{\beta_{n-2}} + \dots + \frac{\alpha_1}{\beta_1}\right)]] \int_0^t f(\tau) \tau^{\frac{\alpha_1}{\beta_1} + \frac{\alpha_2}{\beta_2} + \dots + \frac{\alpha_n}{\beta_n} - 1} d\tau \quad (9)$$

4. Extensions

We now consider for $r = 1, 2, \dots, n$ and $0 \leq s \leq n - r$, the integral

$$J_{r,s} = \int \int \dots \int_R f(t_r + t_{r+1} + \dots + t_{r+s}) t_1^{\alpha_1-1} t_2^{\alpha_2-1} \dots t_n^{\alpha_n-1} dt_1 dt_2 \dots dt_n$$

Where R is the same region as described in (3). The function $f(\cdot)$ and the parameters $\alpha_1, \alpha_2, \dots, \alpha_n$ are as defined before. It is possible to reduce $J_{1,n-2}$ and $J_{n,0}$ to single integrals although a simple reduction does not appear to exist for other values of r and s

a. The Integral $J_{1,n-2}$

We first show that a double integral of the form

$$H(a, b) = \int_0^b \int_0^a f[\tau(1-v)A][\tau(1-v)]^{\alpha-1} (\tau v)^{\beta-1} \tau d\tau dv$$

where $0 \leq a, b \leq 1$; $A > 0$; $\alpha, \beta > 0$ is reducible to single integrals. Making the change in variable $\tau(1-v) = \theta$ and $(1-v) = u$, we get

$$H(a, b) = \int_{1-b}^1 \int_0^{au} f(A\theta)\theta^{\alpha+\beta-1} (1-u)^{\beta-1} u^{-\beta-1} d\theta du$$

Interchanging the order of integration yields

$$\begin{aligned} H(a, b) &= \int_0^a f(A\theta)\theta^{\alpha+\beta-1} \int_{\frac{\theta}{a}}^1 (1-u)^{\beta-1} u^{-\beta-1} du d\theta \\ &- \int_0^{a(1-b)} f(A\theta)\theta^{\alpha+\beta-1} \int_{\frac{\theta}{a}}^{1-b} (1-u)^{\beta-1} u^{-\beta-1} du d\theta \end{aligned}$$

Making the change in variable $t = (1-u)/u$ we obtain finally

$$\begin{aligned} H(a, b) &= \frac{1}{\beta} \int_0^a f(A\theta)\theta^{\alpha+\beta-1} \left(\frac{a}{\theta} - 1\right)^{\beta} d\theta \\ &- \frac{1}{\beta} \int_0^{a(1-b)} f(A\theta)\theta^{\alpha+\beta-1} \left[\left(\frac{a}{\theta} - 1\right)^{\beta} - \left(\frac{b}{1-b} - 1\right)^{\beta}\right] d\theta \end{aligned}$$

Now

$$J_{1,n-2} = \int \int \cdots \int_R f(t_1 + t_2 + \cdots + t_{n-1}) t_1^{\alpha_1-1} t_2^{\alpha_2-1} \cdots t_n^{\alpha_n-1} dt_1 dt_2 \cdots dt_n$$

To evaluate $J_{1,n-2}$ we perform the same sequence of variable transformation as in the evaluation of J_n , and obtain

$$J_{1,n-2} = [B_{\frac{a_1}{1+a_1}}(\alpha_2, \alpha_1)] [B_{\frac{a_2}{1+a_2}}(\alpha_3, \alpha_2 + \alpha_1)] \cdots [B_{\frac{a_{n-2}}{1+a_{n-2}}}(\alpha_{n-1}, \alpha_{n-2} + \alpha_{n-3} + \cdots + \alpha_1)] \\ \int_0^{\frac{a_{n-1}}{1+a_{n-1}}} \int_0^1 f[\tau(1 - u_{n-1})] [\tau(1 - u_{n-1})]^{\alpha_1 + \alpha_2 + \cdots + \alpha_{n-1} - 1} (\tau u_{n-1})^{\alpha_n - 1} \tau d\tau du_{n-1}$$

This double integral is of the same form as $H(a, b)$ with $A = 1$, $a = 1$,

$b = a_{n-1}/(1 + a_{n-1})$, $\alpha = \alpha_1 + \alpha_2 + \cdots + \alpha_{n-1}$, $\beta = \alpha_n$. Hence

$$J_{1,n-2} = [B_{\frac{a_1}{1+a_1}}(\alpha_2, \alpha_1)] [B_{\frac{a_2}{1+a_2}}(\alpha_3, \alpha_2 + \alpha_1)] \cdots [B_{\frac{a_{n-2}}{1+a_{n-2}}}(\alpha_{n-1}, \alpha_{n-2} + \alpha_{n-3} + \cdots + \alpha_1)] \\ \left\{ \frac{1}{\alpha_n} \int_0^1 f(\theta) \theta^{\alpha_1 + \alpha_2 + \cdots + \alpha_{n-1} - 1} \left(\frac{1}{\theta} - 1\right)^{\alpha_n} d\theta \right. \\ \left. - \frac{1}{\alpha_n} \int_0^{\frac{1}{1+a_{n-1}}} f(\theta) \theta^{\alpha_1 + \alpha_2 + \cdots + \alpha_{n-1} - 1} \left[\left(\frac{1}{\theta} - 1\right)^{\alpha_n} - (a_{n-1})^{\alpha_n}\right] d\theta \right\}$$

and in final form

$$J_{1,n-2} = [B_{\frac{a_1}{1+a_1}}(\alpha_2, \alpha_1)] [B_{\frac{a_2}{1+a_2}}(\alpha_3, \alpha_2 + \alpha_1)] \cdots [B_{\frac{a_{n-2}}{1+a_{n-2}}}(\alpha_{n-1}, \alpha_{n-2} + \alpha_{n-3} + \alpha_{n-3} + \cdots + \alpha_1)]$$

$$\left\{ \left[\int_0^1 - \int_0^{\frac{1}{1+a_{n-1}}} \right] f(\theta) \theta^{\alpha_1 + \alpha_2 + \dots + \alpha_{n-1} - 1} (1 - \theta)^{\alpha_n} d\theta \right. \\ \left. + \frac{(a_{n-1})^{\alpha_n}}{\alpha_n} \int_0^{\frac{1}{1+a_{n-1}}} f(\theta) \theta^{\alpha_1 + \alpha_2 + \dots + \alpha_n - 1} d\theta \right\}$$

b. The Integral $J_{n,0}$

We consider here a double integral of the form

$$\tilde{K}(a, b) = \int_0^b \int_0^a f(\tau v A) [\tau(1-v)]^{\alpha-1} (\tau v)^{\beta-1} \tau d\tau dv$$

where $0 \leq a, b \leq 1$; $A > 0$; $\alpha, \beta > 0$. Making the change in variable $\tau v = \theta$ and $v = u$, we get

$$\begin{aligned} \tilde{K}(a, b) &= \int_0^b \int_0^{au} f(A\theta) \theta^{\alpha+\beta-1} (1-u)^{\alpha-1} u^{-\alpha-1} d\theta du \\ &= \int_0^{ab} f(A\theta) \theta^{\alpha+\beta-1} \int_{\frac{\theta}{a}}^b (1-u)^{\alpha-1} u^{-\alpha-1} du d\theta \\ &= \frac{1}{\alpha} \int_0^{ab} f(A\theta) \theta^{\alpha+\beta-1} \left[-\left(\frac{1}{b} - 1\right)^{\alpha} + \left(\frac{a}{\theta} - 1\right)^{\alpha} \right] d\theta \\ &= -\frac{1}{\alpha} \left(\frac{1}{b} - 1\right)^{\alpha} \int_0^{ab} f(A\theta) \theta^{\alpha+\beta-1} d\theta + \frac{1}{\alpha} \int_0^{ab} f(A\theta) \theta^{\beta-1} (a - \theta)^{\alpha} d\theta \end{aligned}$$

Now

$$J_{n,0} = \int \int \dots \int_R f(t_n) t_1^{\alpha_1-1} t_2^{\alpha_2-1} \dots t_n^{\alpha_n-1} dt_1 dt_2 \dots dt_n$$

Again, we perform the same sequence of variable transformation as in the evaluation of J_n and we obtain

$$\begin{aligned} J_{n,0} &= \left[B_{a_1}^{(\alpha_2, \alpha_1)} \right] \left[B_{a_2}^{(\alpha_3, \alpha_2 + \alpha_1)} \right] \dots \left[B_{a_{n-2}}^{(\alpha_{n-1}, \alpha_{n-2} + \alpha_{n-3} + \dots + \alpha_1)} \right] \\ &\quad \int_0^{\frac{a_{n-1}}{1+a_{n-1}}} \int_0^1 f(\tau u_{n-1}) \\ &\quad [\tau(1 - u_{n-1})]^{\alpha_1 + \alpha_2 + \dots + \alpha_{n-1} - 1} (\tau u_{n-1})^{\alpha_n - 1} \tau d\tau du_{n-1} \end{aligned}$$

This double integral is of the same form as $\tilde{K}(a, b)$ with $A = 1$, $a = 1$,
 $b = a_{n-1}/(1 + a_{n-1})$, $\alpha = \alpha_1 + \alpha_2 + \dots + \alpha_{n-1}$ and $\beta = \alpha_n$. Hence

$$J_{n,0} = [B \frac{a_1}{1+a_1} (\alpha_2, \alpha_1)] [B \frac{a_2}{1+a_2} (\alpha_3, \alpha_2 + \alpha_1)] \dots [B \frac{a_{n-2}}{1+a_{n-2}} (\alpha_{n-1}, \alpha_{n-2} + \alpha_{n-3} + \dots + \alpha_1)].$$

$$\frac{1}{\alpha_1 + \alpha_2 + \dots + \alpha_{n-1}} \left[\int_0^{\frac{a_{n-1}}{1+a_{n-1}}} f(\theta) \theta^{\alpha_{n-1}-1} (1-\theta)^{\alpha_1+\alpha_2+\dots+\alpha_{n-1}} d\theta \right.$$

$$\left. - \left(\frac{1}{a_{n-1}} \right)^{\alpha_1+\alpha_2+\dots+\alpha_{n-1}} \int_0^{\frac{a_{n-1}}{1+a_{n-1}}} f(\theta) \theta^{\alpha_1+\alpha_2+\dots+\alpha_{n-1}-1} d\theta \right]$$

5. A Still Further Extension.

Another multiple integral which is reducible to a single integral is the following

$$\int \int \dots \int_R f(t_1 + t_2 + \dots + t_n) t_1^{\alpha_1-1} t_2^{\alpha_2-1} \dots t_n^{\alpha_n-1} \\ (t_1 + t_2)^{\beta_2} (t_1 + t_2 + t_3)^{\beta_3} \dots (t_1 + t_2 + \dots + t_{n-1})^{\beta_{n-1}} \\ dt_1 dt_2 \dots dt_n$$

where $\beta_i \geq 0$, ($i = 2, 3, \dots, n-1$). The reduction process is similar to that used for (4) and will not be repeated here.

6. Applications to Statistics

Statistical distributions of the multivariate type classified by functional form have received little attention. For univariate classes, the Pearson's type (e.g. see [6]) are well known. For the multivariate case, Lord [14] and Box and Hunter [3] consider the radical or spherical distributions where the joint probability density functions of the sequence of random variables

$\{X_1, X_2, \dots, X_n\}$ has the form

$$\phi_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = A f(x_1^2 + x_2^2 + \dots + x_n^2)$$

where A is a constant. Kelker [11] studies linear transformations of spherically distributed variables, while Baldessari [2] considers density functions of the form

$$\phi_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = A f(x_1 + x_2 + \dots + x_n)$$

where $\sum_{j=1}^{j=n} (x_j - a)^2 \leq r^2$, and of the form

$$\phi_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) = A f[(x_1 - a_1)^2 + (x_2 - a_2)^2 + \dots + (x_n - a_n)^2]$$

where $\sum_{j=1}^{j=n} (x_j - a_j)^2 \leq r^2$. Finally, the so-called elliptically symmetric

distributions are studied by McGraw and Wagner [15].

Power transformations of the type $X = Y^\lambda$ have been used in Dirichlet's distributions (see e.g. Johnston and Kotz [9]) and in the form $X = aY^\lambda - (1 - Y)^\lambda$ are known as the Tukey lambda transformation [8][10]. Generalizations to the standard Dirichlet distribution were considered by Johnston and Kotz [9].

We consider presently some generalizations of the previously investigated classes of statistical distributions. In particular, we study some properties of a class of multivariate distributions defined by a functional form in

which the variables may be subject to a power transformation.

Let then A be a positive constant and $f(x)$ be a continuous nonnegative function defined everywhere in the interval $0 \leq x < \infty$ with the property that for $\gamma_i > 0$, $i = 1, 2, \dots, n$, the improper integral

$$\int_0^{\infty} f(\tau) \tau^{\gamma_1 + \gamma_2 + \dots + \gamma_n - 1} d\tau$$

exists and converges absolutely.

Let $\{X_1, X_2, \dots, X_n\}$ be an n -dimensional non-negative random variable whose P.D.F. $\phi_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n)$ is equal to

$$A f\left[\left(\frac{x_1}{q_1}\right)^{\beta_1} + \left(\frac{x_2}{q_2}\right)^{\beta_2} + \dots + \left(\frac{x_n}{q_n}\right)^{\beta_n}\right] x_1^{\alpha_1 - 1} x_2^{\alpha_2 - 1} \dots x_n^{\alpha_n - 1} \quad (10)$$

over the region $\{(x_1, x_2, \dots, x_n) : 0 \leq x_i < \infty, i = 1, 2, \dots, n\}$, and is zero elsewhere, and let for $i = 1, 2, \dots, n$ the quantities $\alpha_i > 0$, $\beta_i > 0$, $q_i > 0$ be uniquely defined.

a. The Constant A

The quantity A is given by the normalizing conditions

$$\frac{1}{A} = \int_0^{\infty} \int_0^{\infty} \dots \int_0^{\infty} f\left[\left(\frac{x_1}{q_1}\right)^{\beta_1} + \left(\frac{x_2}{q_2}\right)^{\beta_2} + \dots + \left(\frac{x_n}{q_n}\right)^{\beta_n}\right] x_1^{\alpha_1 - 1} x_2^{\alpha_2 - 1} \dots x_n^{\alpha_n - 1} dx_1 dx_2 \dots dx_n$$

The multiple integral is of the same form as (8) with $t = \infty$ and $a_i = \infty$, $i = 1, 2, \dots, n$. Hence

$$\frac{1}{A} = \frac{q_1^{\alpha_1} q_2^{\alpha_2} \dots q_n^{\alpha_n}}{\beta_1 \beta_2 \dots \beta_n} \frac{\Gamma\left(\frac{\alpha_1}{\beta_1}\right) \Gamma\left(\frac{\alpha_2}{\beta_2}\right) \dots \Gamma\left(\frac{\alpha_n}{\beta_n}\right)}{\Gamma\left(\frac{\alpha_1}{\beta_1} + \frac{\alpha_2}{\beta_2} + \dots + \frac{\alpha_n}{\beta_n}\right)} \int_0^{\infty} f(\tau) \tau^{\frac{\alpha_1}{\beta_1} + \frac{\alpha_2}{\beta_2} + \dots + \frac{\alpha_n}{\beta_n} - 1} d\tau \quad (11)$$

b. Mixed Moments

The mixed moments about the origin if they exist, are defined for

$r_1 = 1, 2, \dots, (i = 1, 2, \dots, n)$ by the expression

$$\begin{aligned} \mu'_{r_1 r_2 \dots r_n} &= \int_0^\infty \int_0^\infty \dots \int_0^\infty x_1^{r_1} x_2^{r_2} \dots x_n^{r_n} \phi_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) dx_1, dx_2, \dots, dx_n \\ &= A \frac{q_1^{\alpha_1+r_1} q_2^{\alpha_2+r_2} \dots q_n^{\alpha_n+r_n}}{\beta_1 \beta_2 \dots \beta_n} \frac{\Gamma(\frac{\alpha_1+r_1}{\beta_1}) \Gamma(\frac{\alpha_2+r_2}{\beta_2}) \dots \Gamma(\frac{\alpha_n+r_n}{\beta_n})}{\Gamma(\frac{\alpha_1+r_1}{\beta_1} + \frac{\alpha_2+r_2}{\beta_2} + \dots + \frac{\alpha_n+r_n}{\beta_n})} \\ &\quad \int_0^\infty f(\tau) \tau^{\frac{\alpha_1+r_1}{\beta_1} + \frac{\alpha_2+r_2}{\beta_2} + \dots + \frac{\alpha_n+r_n}{\beta_n} - 1} d\tau \end{aligned}$$

c. Marginal Distributions

The marginal PDF $\phi_{X_1}(\cdot)$ of X_1 is given by ($0 \leq x_1 < \beta$)

$$\begin{aligned} \phi_{X_1}(x_1) &= \int_0^\infty \int_0^\infty \dots \int_0^\infty \phi_{X_1, X_2, \dots, X_n}(x_1, x_2, \dots, x_n) dx_2 dx_3 \dots dx_{i-1} dx_{i+1} \dots dx_n \\ &= A \frac{q_1^{\alpha_1} q_2^{\alpha_2} \dots q_{i-1}^{\alpha_{i-1}} q_{i+1}^{\alpha_{i+1}} \dots q_n^{\alpha_n}}{\beta_1 \beta_2 \dots \beta_{i-1} \beta_{i+1} \dots \beta_n} \\ &\quad \int_0^\infty f\left[\tau + \left(\frac{x_1}{q_1}\right)^{\beta_1}\right] \tau^{\frac{\alpha_1}{\beta_1} + \frac{\alpha_2}{\beta_2} + \frac{\alpha_{i-1}}{\beta_{i-1}} + \frac{\alpha_{i+1}}{\beta_{i+1}} + \dots + \frac{\alpha_n}{\beta_n} - 1} d\tau \end{aligned}$$

By a suitable change in the order of integration, setting $y = \tau + (x_1/q_1)^{\beta_1}$, and using the convolution theorem, one can verify that $\int_0^\infty \phi_{X_1}(x_1) = 1$. In a similar fashion, it is possible to show that the joint marginal P.D.F. of $\{X_1, X_2, \dots, X_k\}$ ($k \leq n$) is given by

$$\phi_{X_1, X_2, \dots, X_k}(x_1, x_2, \dots, x_k) = A \frac{q_{k+1}^{a_{k+1}} q_{k+2}^{a_{k+2}} \dots q_n^{a_n}}{\beta_{k+1} \beta_{k+2} \dots \beta_n} x_1^{a_1-1} x_2^{a_2-1} \dots x_k^{a_k-1} \\ \int_0^\infty f\left[\left(\frac{x_1}{q_1}\right)^{\beta_1} + \left(\frac{x_2}{q_2}\right)^{\beta_2} + \dots + \left(\frac{x_n}{q_n}\right)^{\beta_n} + \tau\right] \tau^{\frac{a_{k+1}}{\beta_{k+1}} + \frac{a_{k+2}}{\beta_{k+2}} + \dots + \frac{a_n}{\beta_n} - 1} d\tau$$

The conditional distributions can then readily be obtained. We presently establish the following two theorems

d. Theorem 1

The $(n - 1)$ random variables $\{Y_1, Y_2, \dots, Y_{n-1}\}$ where

$$Y_1 = \frac{\frac{x_1}{q_1}^{\beta_1}}{\frac{x_1}{q_1}^{\beta_1} + \frac{x_2}{q_2}^{\beta_2} + \dots + \frac{x_n}{q_n}^{\beta_n}}, \quad i = 1, 2, \dots, n - 1$$

have a joint probability density function which is of the Dirichlet type inside the simplex

$$\mathcal{J} = \{(y_1, y_2, \dots, y_{n-1}) : y_1 + y_2 + \dots + y_{n-1} \leq 1, y_1 \geq 0, \\ y_2 \geq 0, \dots, y_{n-1} \geq 0\}$$

and which is zero elsewhere.

Proof:

$$\text{Let } Y_n = \frac{x_1}{q_1}^{\beta_1} + \dots + \frac{x_n}{q_n}^{\beta_n}$$

$$\text{Then } \frac{x_1}{q_1}^{\beta_1} = Y_1 Y_n, \quad i = 1, 2, \dots, n - 1$$

$$\text{and } \frac{x_n}{q_n}^{\beta_n} = (1 - Y_1 - Y_2 - \dots - Y_{n-1}) Y_n$$

This particular transformation maps the region $\{(x_1, x_2, \dots, x_n) : 0 \leq x_i < \infty, i = 1, 2, \dots, n\}$ onto the region $C = \{(y_1, y_2, \dots, y_n) : y_1 + y_2 + \dots + y_{n-1} \leq 1, y_i \geq 0 \text{ for } i = 1, 2, \dots, n-1, 0 \leq y_n < \infty\}$. The corresponding inverse functions for this one to one transformation is

$$x_i = q_i (y_i y_n)^{1/\beta_i}, \quad i = 1, 2, \dots, n-1$$

$$x_n = q_n [(1 - y_1 - y_2 - \dots - y_{n-1}) y_n]^{1/\beta_n}$$

The Jacobian of the transformation is

$$J = \frac{\partial(x_1, x_2, \dots, x_n)}{\partial(y_1, y_2, \dots, y_n)}$$

$$= \begin{vmatrix} \frac{\partial x_1}{\partial y_1} & \frac{\partial x_1}{\partial y_2} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \frac{\partial x_n}{\partial y_2} & \dots & \frac{\partial x_n}{\partial y_n} \end{vmatrix}$$

Now for $i = 1, 2, \dots, n$, we have

$$\frac{\partial x_i}{\partial y_i} = \frac{q_i}{\beta_i} (y_i y_n)^{\frac{1}{\beta_i} - 1} y_n$$

$$\frac{\partial x_i}{\partial y_n} = \frac{q_i}{\beta_i} (y_i y_n)^{\frac{1}{\beta_i} - 1} y_i$$

$$\frac{\partial x_i}{\partial y_j} = 0 \quad \text{for } j \neq i, n$$

Also for $i = 1, 2, \dots, n-1$

$$\frac{\partial x_n}{\partial y_i} = \frac{q_n}{\beta_n} (-y_n) [(1 - y_1 - y_2 - \dots - y_{n-1}) y_n]^{\frac{1}{\beta_n} - 1}$$

and
$$\frac{\partial x_n}{\partial y_n} = \frac{q_n}{\beta_n} (1 - y_1 - y_2 - \dots - y_{n-1}) [(1 - y_1 - y_2 - \dots - y_{n-1}) y_n]^{\frac{1}{\beta_n} - 1}$$

The value of the Jacobian is thus

$$J = \frac{q_1}{\beta_1} (y_1 y_n)^{\frac{1}{\beta_1} - 1} \cdot \frac{q_2}{\beta_2} (y_2 y_n)^{\frac{1}{\beta_2} - 1} \dots \frac{q_{n-1}}{\beta_{n-1}} (y_{n-1} y_n)^{\frac{1}{\beta_{n-1}} - 1} \cdot \frac{q_n}{\beta_n} [(1 - y_1 - y_2 - \dots - y_{n-1}) y_n]^{\frac{1}{\beta_n} - 1}$$

$$\begin{vmatrix} y_n & 0 & \dots & y_1 \\ 0 & y_n & \dots & y_2 \\ \vdots & \vdots & \dots & \vdots \\ -y_n & -y_n & \dots & (1 - y_1 - y_2 - \dots - y_{n-1}) \end{vmatrix}$$

Noting that the value of the determinant is y_n^{n-1} , we have

$$J = \frac{q_1 q_2 \dots q_n}{\beta_1 \beta_2 \dots \beta_n} y_1^{\frac{1}{\beta_1} - 1} y_2^{\frac{1}{\beta_2} - 1} \dots y_{n-1}^{\frac{1}{\beta_{n-1}} - 1} y_n^{\frac{1}{\beta_1} + \frac{1}{\beta_2} + \dots + \frac{1}{\beta_n} - 1} (1 - y_1 - y_2 - \dots - y_{n-1})^{\frac{1}{\beta_n} - 1}$$

Hence, the joint P.D.F. of $\{Y_1, Y_2, \dots, Y_{n-1}, Y_n\}$ is for $(y_1, y_2, \dots, y_n) \in C$

$$A f(y_n) q_1^{\alpha_1 - 1} (y_1 y_n)^{\frac{\alpha_1 - 1}{\beta_1}} \cdot q_2^{\alpha_2 - 1} (y_2 y_n)^{\frac{\alpha_2 - 1}{\beta_2}} \dots q_{n-1}^{\alpha_{n-1} - 1} (y_{n-1} y_n)^{\frac{\alpha_{n-1} - 1}{\beta_{n-1}}} q_n^{\alpha_n - 1} [(1 - y_1 - y_2 - \dots - y_{n-1}) y_n]^{\frac{\alpha_n - 1}{\beta_n}} \cdot |J|$$

$$= A \frac{q_1^{\alpha_1} q_2^{\alpha_2} \dots q_n^{\alpha_n}}{\beta_1 \beta_2 \dots \beta_n} y_1^{\frac{\alpha_1}{\beta_1} - 1} y_2^{\frac{\alpha_2}{\beta_2} - 1} \dots y_{n-1}^{\frac{\alpha_{n-1}}{\beta_{n-1}} - 1} \cdot (1 - y_1 - y_2 - \dots - y_{n-1})^{\frac{\alpha_n}{\beta_n} - 1} \\ \cdot f(y_n) y_n^{\frac{\alpha_1}{\beta_1} + \frac{\alpha_2}{\beta_2} + \dots + \frac{\alpha_n}{\beta_n} - 1}$$

and is zero elsewhere. Using the computed value of the constant A in (11) we obtain for the joint P.D.F. of $\{Y_1, Y_2, \dots, Y_{n-1}\}$ inside the simplex Δ to be

$$\frac{\Gamma(\frac{\alpha_1}{\beta_1} + \frac{\alpha_2}{\beta_2} + \dots + \frac{\alpha_n}{\beta_n})}{\Gamma(\frac{\alpha_1}{\beta_1}) \Gamma(\frac{\alpha_2}{\beta_2}) \dots \Gamma(\frac{\alpha_n}{\beta_n})} y_1^{\frac{\alpha_1}{\beta_1} - 1} y_2^{\frac{\alpha_2}{\beta_2} - 1} \dots y_{n-1}^{\frac{\alpha_{n-1}}{\beta_{n-1}} - 1} \\ \cdot (1 - y_1 - y_2 - \dots - y_{n-1})^{\frac{\alpha_n}{\beta_n} - 1}$$

and to be zero elsewhere.

e. Theorem 2

The n random variables $\{Y_1, Y_2, \dots, Y_n\}$ where

$$Y_i = \frac{\frac{x_{i+1}}{(q_{i+1})^{\beta_{i+1}}}}{\frac{x_1}{(q_1)^{\beta_1}} + \frac{x_2}{(q_2)^{\beta_2}} + \dots + \frac{x_i}{(q_i)^{\beta_i}}} \quad i = 1, 2, \dots, n-1$$

$$Y_n = \frac{x_1}{(q_1)^{\beta_1}} + \frac{x_2}{(q_2)^{\beta_2}} + \dots + \frac{x_n}{(q_n)^{\beta_n}}$$

are independently distributed. The marginal P.D.F. of Y_i 's are

$$\phi_{Y_i}(y_i) = \begin{cases} A_i \frac{\frac{\alpha_{i+1}}{\beta_{i+1}} - 1}{(1 + y_i)^{\frac{\alpha_1}{\beta_1} + \frac{\alpha_2}{\beta_2} + \dots + \frac{\alpha_{i+1}}{\beta_{i+1}}}}, & 0 < y_i < \infty \\ 0, & \text{otherwise} \end{cases} \quad (i = 1, 2, \dots, n-1)$$

$$\phi_{Y_n}(y_n) = \begin{cases} A_n \frac{\frac{\alpha_1}{\beta_1} + \frac{\alpha_2}{\beta_2} + \dots + \frac{\alpha_n}{\beta_n} - 1}{f(y_n) y_n}, & 0 < y_n < \infty \\ 0, & \text{otherwise} \end{cases}$$

where the A_i 's are normalizing constants.

Proof:

The joint distribution function of $\{Y_1, Y_2, \dots, Y_n\}$ is given by

$$P\{Y_1 \leq y_1, Y_2 \leq y_2, \dots, Y_n \leq y_n\} =$$

$$P\left\{\frac{x_2}{(q_2)^{\beta_2}} \leq y_1 \frac{x_1}{(q_1)^{\beta_1}}, \frac{x_3}{(q_3)^{\beta_3}} \leq y_2 \left[\frac{x_1}{(q_1)^{\beta_1}} + \frac{x_2}{(q_2)^{\beta_2}}\right], \dots,\right.$$

$$\left. \frac{x_n}{(q_n)^{\beta_n}} \leq y_{n-1} \sum_{i=1}^{n-1} \frac{x_i}{(q_i)^{\beta_i}}, \sum_{i=1}^n \frac{x_i}{(q_i)^{\beta_i}} \leq y_n\right\}$$

$$= A \int \int \dots \int_{\tilde{T}} f\left[\left(\frac{x_1}{q_1}\right)^{\beta_1} + \left(\frac{x_2}{q_2}\right)^{\beta_2} + \dots + \left(\frac{x_n}{q_n}\right)^{\beta_n}\right] x_1^{\alpha_1-1} x_2^{\alpha_2-1} \dots x_n^{\alpha_n-1} dx_1 dx_2 \dots dx_n$$

where \tilde{T} is the region of integration defined by the set of points

$$\tilde{T} = \{(x_1, x_2, \dots, x_n) : x_i \geq 0, \sum_{j=1}^n \left(\frac{x_j}{q_j}\right)^{\beta_j} \leq y_n, \left(\frac{x_{i+1}}{q_{i+1}}\right)^{\beta_{i+1}} \leq y_i \sum_{j=1}^n \left(\frac{x_j}{q_j}\right)^{\beta_j}; i = 1, 2, \dots, n\}$$

Thus, from (9)

$$P\{Y_1 \leq y_1, Y_2 \leq y_2, \dots, Y_n \leq y_n\} = A \frac{q_1^{\alpha_1} q_2^{\alpha_2} \dots q_n^{\alpha_n}}{\beta_1 \beta_2 \dots \beta_n} \left[B \frac{y_1}{1+y_1} \left(\frac{\alpha_2}{\beta_2}, \frac{\alpha_1}{\beta_1} \right) \right] \cdot \left[B \frac{y_2}{1+y_2} \left(\frac{\alpha_3}{\beta_3}, \frac{\alpha_2}{\beta_2} + \frac{\alpha_1}{\beta_1} \right) \right] \dots \left[B \frac{y_{n-1}}{1+y_{n-1}} \left(\frac{\alpha_n}{\beta_n}, \frac{\alpha_{n-1}}{\beta_{n-1}} + \frac{\alpha_{n-2}}{\beta_{n-2}} + \dots + \frac{\alpha_1}{\beta_1} \right) \right] \cdot \int_0^{y_n} f(\tau) \tau^{\frac{\alpha_1}{\beta_1} + \frac{\alpha_2}{\beta_2} + \dots + \frac{\alpha_n}{\beta_n} - 1} dy_n$$

which establishes the independence of Y_1, Y_2, \dots, Y_n . The distribution function of Y_i is $(0 \leq y_i < \infty) (i = 1, 2, \dots, n-1)$

$$A_i B \frac{y_i}{1+y_i} \left(\frac{\alpha_{i+1}}{\beta_{i+1}}, \frac{\alpha_i}{\beta_i} + \frac{\alpha_{i-1}}{\beta_{i-1}} + \dots + \frac{\alpha_1}{\beta_1} \right) = A_i \int_0^{y_i} \frac{y}{1+y} \frac{\alpha_{i+1}}{\beta_{i+1}} - 1 (1-y)^{\frac{\alpha_i}{\beta_i} + \frac{\alpha_{i-1}}{\beta_{i-1}} + \dots + \frac{\alpha_1}{\beta_1} - 1} dy$$

where
$$A_1 = \frac{1}{B\left(\frac{\alpha_1}{\beta_1} + \frac{\alpha_2}{\beta_2} + \dots + \frac{\alpha_i}{\beta_i}, \frac{\alpha_{i+1}}{\beta_{i+1}}\right)}$$

Hence the P.D.F. of Y_1 ($i = 1, 2, \dots, n-1$) is for $0 < y_1 < \infty$

$$\phi_{Y_1}(y_1) = A_1 \frac{y_1^{\frac{\alpha_{i+1}}{\beta_{i+1}} - 1}}{(1 + y_1)^{\frac{\alpha_1}{\beta_1} + \frac{\alpha_2}{\beta_2} + \dots + \frac{\alpha_{i+1}}{\beta_{i+1}}}}$$

which can be recognized as a beta distribution of the second kind. The P.D.F. of Y_n is ($0 \leq y_n < \infty$)

$$\phi_{Y_n}(y_n) = A_n f(y_n) y_n^{\frac{\alpha_1}{\beta_1} + \frac{\alpha_2}{\beta_2} + \dots + \frac{\alpha_n}{\beta_n} - 1}$$

where
$$\frac{1}{A_n} = \int_0^\infty f(\tau) \tau^{\frac{\alpha_1}{\beta_1} + \frac{\alpha_2}{\beta_2} + \dots + \frac{\alpha_n}{\beta_n} - 1} d\tau$$

Theorems 1 and 2 are generalizations of some well known results in statistics (see e.g. Wilks [19] and Aitchison [1]).

References

- [1] J. Aitchison, Inverse Distributions and Independent Gamma-Distributed Products of Random Variables, *Biometrika*, 50 (1963) pp. 505-509.
- [2] B. Baldessari, Osservazioni Sulle Funzioni Caratteristiche Di Alcune Classi Di V. C. n-dimensionali. Pubblicazioni della Seconda Serie dell' Istituto di Calcolo delle Probabilita dell' Universita di Roma, 49 (1965) pp. 1-11.
- [3] G. E. P. Box and J. S. Hunter, Multi-Factor Experimental Designs for Exploring Response Surfaces, *Ann. Math Stat.* 28 (1957); pp. 195-241.
- [4] J. L. Coolidge, *A History of Geometrical Methods*, Oxford University Press, London, 1940.
- [5] J. Edwards, *A Treatise on the Integral Calculus*, Vol. 2. Macmillan, N. Y., 1922.
- [6] W. P. Elderton, *Frequency Curves and Correlation*, 3rd ed. Cambridge, 1938.
- [7] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series and Products*, Academic Press, N.Y., 1965.
- [8] C. Hastings, F. Mosteller, J. W. Tukey and C. P. Winsor, Low Moments for Small Samples: A Comparative Study of Order Statistics, *Ann. Math. Stat.*, 18 (1947); pp. 413-426.

- [9] N. L. Johnson and S. Kotz, Distributions in Statistics: Continuous Multivariate Distributions, J. Wiley & Sons, N. Y., 1972.
- [10] B. L. Joiner and J. R. Rosenblatt, Some Properties of Range in Samples from Tukey's Symmetric Lambda Distributions, Jour. Amer. Stat. Assn., 66 (1971) pp. 992-995.
- [11] D. Kelker, Distribution Theory of Spherical Distributions and a Location-Scale Parameter Generalization, Sankhya, Series A, 32 (1970) pp. 419-430.
- [12] M. G. Kendall and A. Stuart, The Advanced Theory of Statistics, Vol. 1, Charles Griffi & Co. Ltd., London, 1969.
- [13] M. S. Klamkin, Extensions of Dirichlet's Multiple Integral, SIAM J. Math. Anal. 2 (1971) pp. 467-469.
- [14] R. D. Lord, The Use of Hankel Transforms in Statistics, I, General Theory and Examples, Biometrika, 41 (1954); pp. 44-55.
- [15] D. K. McGraw and J. F. Wagner, Elliptically Symmetric Distributions, IEEE Transactions on Information Theory, 14 (1968), pp. 110-120.
- [16] B. D. Sivazlian, The Generalized Dirichlet's Multiple Integral, SIAM Review, 11 (1969) pp. 285-288.
- [17] _____, A Class of Multiple Integrals, SIAM J. Math. Anal. 2 (1971) pp. 72-75.

- [18] E. T. Whittaker and G. N. Watson, A Course of Modern Analysis, Cambridge University Press, London, 1927.
- [19] S. S. Wilks, Mathematical Statistics, J. Wiley & Sons, N. Y., 1962.